

Aminosteril infant 10%

Asam Amino, Taurine

Composition

1000 ml contains:

L-leucine	13.00	g
L-isoleucine	8.00	g
L-lysine acetate	12.00	g
= L-lysine 8.51 g		
L-methionine	3.12	g
L-phenylalanine	3.75	g
L-threonine	4.40	g
L-tryptophan	2.01	g
L-valine	9.00	g
Arginine	7.50	g
L-histidine	4.76	g
Glycine	4.15	g
L-taurine	0.40	g
L-serine	7.67	g
L-alanine	9.30	g
L-proline	9.71	g
N-acetyl-L-tyrosine	5.176	g
= L-tyrosine 4.20 g		
N-acetyl-L-cysteine	0.70	g
= L-cysteine 0.52 g		
L-malic acid	2.62	g
Total amino acids:	100 g/l	
Total nitrogen content:	14.9 g/l	
Titration acidity:	25 - 45 mmol NaOH/l	
pH-value:	5.5 - 6.0	
Theoretical osmolarity:	885 mosm/l	

Other constituents: Water for injections

DRUG ACTION

Parenteral infusion of amino acids is necessary, when oral or enteral supply with protein is impossible. Especially newborn infants, babies and children require constant provision of protein for proper organ function, developments and growth.

Intravenously given amino acids are proven to be metabolized like those derived from nutrient proteins. The composition of aminosteril infant 10% has been specifically formulated to provide an optimal nitrogen source for nutritional support and therapy for pediatric patient. The solution is design to meet the requirements for protein - building substances of pediatric patients in different disease conditions. The amino acid pattern can be assessed suitable for the maintenance of the nutritional status.

INDICATIONS

Parenteral nutrition for prophylaxis and treatment of protein deficiency in pediatrics when oral food intake is either contraindicated or impossible as in cases like :
Premature birth.

Small - for date syndromes.

Gastro - intestinal diseases and malformations.

Peri operatively in pediatric surgery.

Intensive care of infants following burns and severe injuries.

DOSAGE / POSOLOGY

For intravenous infusion

Max. infusion dosage:

Up to 0.1 amino acids / kg b.w. and h = 1.0 ml / kg b.w. & h.

Max. daily dosage:

When administered during the first days of life, dosage should be restricted to 1 g amino acids per kg body weight per day

1st year of living : 1.5 - 2.5 g amino acids / kg b.w.

2nd year of living : 1.5 g amino acids / kg b.w.

3rd to 5th year of living : 1.5 g amino acids / kg b.w.

6th to 10th year of living : 1.0 g amino acids / kg b.w.

11th to 14th year of living : 1.0 g amino acids / kg b.w.

The solution is administered as long as Parenteral nutrition is required.

SPECIAL WARNING / PRECAUTIONS

Do not use Aminosteril Infant 10% after expiry date. Use only clear solution and undamaged containers. Keep drugs out of the reach of children.

Administration of amino acid solution may cause folate deficiency and supplementary folic acid should be administered.

In order to ensure optimum metabolic utilization of the amino acids administered, therapy should be adequately supported by an energy supply of 35 kcal (145 KJ) per gram of amino acids.

Over dosage or a too rapid infusion may manifest in the form of nausea, shivering, vomiting and increased renal amino acids losses.

In such cases, the infusion should be interrupted and eventually continued at a reduced infusion rate.

REMARK

Too rapid infusion may result in renal losses causing amino acids imbalances.

Electrolytes should be administered according to the requirements.

SIDE EFFECT

Local reactions consisting of erythema, phlebitis and thrombosis at the infusions site can occur with peripheral infusion. Generalized flushing, vomiting and fever during infusion of amino acid solution have been reported.

CONTRA INDICATION

Aminosteril infant 10% is contraindicated in patients with inborn errors of amino acids metabolism, hepatic coma and untreated anuria (renal insufficiency), manifest cardiac insufficiency, hyperhydration and hypokalemia.

Patient with insufficient renal or hepatic function require individual dosage.

DRUG INTERACTION

Because of the increased risk of microbiological contaminations and incompatibilities, amino acids solutions should not be mixed with other drugs.

REMARK

The addition of drug may alter the chemical and physical properties of the solution and may therefore give rise to toxic reactions.

Should it nevertheless become necessary to add drugs to Aminosteril Infant 10%, it is imperative to ensure sterility, complete mixing and compatibility. The same applies for the adding of lipid emulsions, trace elements and vitamin for complete parenteral nutrition mixtures.

PRESENTATION

Bottle 100 ml Reg No: DK11681203349E1

Store below 30°C, protect from light.

Shelf life: 2 years

HARUS DENGAN RESEP DOKTER

Manufactured by

Fresenius Kabi Austria, GmbH
Hafnerstrabe 36, A-8055 Craz

For

Fresenius Kabi Deutschland GmbH
61346 Bad Homburg v.d.H
Germany

Imported by

PT. Fresenius Kabi Combiphar
Bandung - Indonesia

 **FRESENIUS
KABI**



M080253/00 ID